

AD-A102 111

SCRIPPS INSTITUTION OF OCEANOGRAPHY LA JOLLA CA

F/6 8/10

ADS REPORT NUMBER 4.(U)

APR 81 S PAZAN, J BYTOW

N00014-80-C-0440

UNCLASSIFIED

SIO-REF.-81-9

NL

1 OF 1  
AD-A  
102111

END  
DATE  
FILMED  
8-81  
DTIC

LEVEL II

12

AD A102111

# SIO REFERENCE SERIES

ADS REPORT NUMBER 4

Issued by NORPAX DATA MANAGEMENT  
S. Pazan, NORPAX Data Manager

DTIC  
ELECTE  
JUL 27 1981  
S D E

DTIC FILE COPY

DISTRIBUTION STATEMENT A  
Approved for public release;  
Distribution Unlimited

April 1981

University of California

SIO Reference 81-9  
Scripps Institution of Oceanography

01 07 11

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER SIO-Ref-81-9/	2. GOVT ACCESSION NO. AD-A102	3. RECIPIENT'S CATALOG NUMBER 111
4. TITLE (and Subtitle) ADS REPORT NUMBER 4.		5. TYPE OF REPORT & PERIOD COVERED Dept. for Nov 77 - Nov 78
7. AUTHOR(s) S. Pazan and J. Bytof		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Scripps Institution of Oceanography La Jolla, CA 92093		8. CONTRACT OR GRANT NUMBER(s) N00014-80-C-0440
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, VA 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 15. SECURITY CLASS. (of this report) Unclassified		12. REPORT DATE April 1981
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release. Distribution unlimited.		13. NUMBER OF PAGES 60
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Since 1976, three reports on oceanographic and meteorological conditions in the North Pacific have been produced by the NORPAX Data Program covering the period from June, 1976 through November, 1977. This is the fourth in the series, and covers the period from December, 1977 through May, 1978. This issue contains: contour maps of monthly meaned Fleet Numerical Oceanographic Central (FNOC) air temperature, sea temperature, wind speed, wind direction, surface vapor pressure, and 700 mb height; NORPAX...		

DD FORM 1473

1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE

S N 0102-LF-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

19-100

CURL OF WIND STRESS CORRIGENDUM TO ADS REPORT NUMBER 4

Curl of wind stress maps in the ADS 4 report are incorrect. These are the corrected maps. The discussion in the Appendix to ADS Report number 4 is correct and pertains to this corrigendum.

*Corrected maps inserted in report.*

*6 Aug 81*

2

NORPAX  
ADS REPORT  
NUMBER 4  
Issued by NORPAX DATA MANAGEMENT  
S. Pazan, J. Bytof  
copyright 1981



The REGENTS of the UNIVERSITY OF CALIFORNIA  
SIO Reference No. 81-9  
April 1981

Accession For		
NTIS	GRA&I	<input checked="checked" type="checkbox"/>
DTIC	TAB	<input type="checkbox"/>
Unannounced		<input type="checkbox"/>
Justification		
By		
Distribution/		
Availability Codes		
Dist	Avail and/or	
	Special	
A		

TABLE OF CONTENTS

INTRODUCTION .....	3
CONTOUR MAPS .....	3
FREE DRIFTER BUOY DISPLACEMENT VECTORS .....	3
FIGURES .....	5
Appendix .....	55



## INTRODUCTION

Since 1976, three reports on oceanographic and meteorological conditions in the North Pacific have been produced by the NORPAX Data Program covering the period from June, 1976 through November, 1977. This is the fourth in the series, and covers the period from December, 1977 through May, 1978. This issue contains: contour maps of monthly meaned Fleet Numerical Oceanographic Central (FNOC) air temperature, sea temperature, wind speed, wind direction, surface vapor pressure, and 700 mb height; NORPAX Data Management calculated wind stress, wind stress curl, wind shear velocity cubed, sensible heat flux, latent heat flux; objectively analyzed TRANSPAC temperatures at discrete depths from White and Bernstein (SIO); monthly ADS buoy drifter displacement vectors from McNally (SIO).

## CONTOUR MAPS

Contour maps of FNOC (Fleet Numerical Oceanographic Central) fields and calculated flux fields are in Figures 2.1-2.11, 3.1-3.11, 4.1-4.11, 5.1-5.11, 6.1-6.11, and 7.1-7.11. TRANSPAC contour maps are in Figures 2.12, 3.12, 4.12, 5.12, 6.12, and 7.12.

The analysis of FNOC fields and NORPAX calculated flux fields are explained in the Appendix to this issue.

XBT's have been regularly dropped from ships of opportunity in the Pacific since 1974. Recovered temperature profile data have been analyzed at Scripps by Bernstein and White and temperature residuals from their anomalies were contoured for 0, 60, 120, 200, 300 and 400 meter depths, by month.

## FREE DRIFTER BUOY DISPLACEMENT VECTORS

Buoy displacement vectors were plotted in Figures 2.13, 3.13, 4.13, 5.13, 6.13, and 7.13. The buoys were drogued at 30 meters.

#### ACKNOWLEDGEMENTS

First, to Anna Moore for her work organizing the figures, xeroxing them using the painstaking Standard Xerox process, we wish to give thanks. Also, to Ted Walker for producing the plots of subsurface temperature fields.






FIGURE 1. 33 x 63 field geographic coverage. The area of the ADS contour maps is as indicated. This map is a polar projection of the northern hemisphere.

WIND SPEED (M/SEC)

DEC 77




FIGURE 2.1 Absolute value of monthly mean vector wind velocities at 19.5 meters. Contour intervals are 1 m/sec.

WIND DIRECTION

DEC

77




FIGURE 2.2 Direction arrows representing directions of monthly mean wind vectors at 19.5 meters. Length of arrow shaft indicates wind speed in m/sec. (See scale above figure).




FIGURE 2.3 Monthly mean wind stress is the mean of 6-hourly wind stress at 10 meters calculated from FNWC wind data. Contour intervals are 0.2 dynes/cm<sup>2</sup>.




FIGURE 2.4 The vertical component of monthly mean wind stress curl is the mean of 6-hourly wind stress curl approximated by finite-differences from 6-hourly wind stresses at 10 meters. Isolines of zero curl are plotted heavily, and contour intervals are  $4.0 \times 10^{-9}$  dynes/cm<sup>3</sup>.

U STAR CUBED (M/SEC) \*\* 3)

DEC 77




FIGURE 2.5 Monthly mean wind shear stress velocity cubed,  $U_*^3$  is the mean of 6-hourly wind shear stress velocity cubed calculated from wind speed at 10 meters. Contour intervals are 0.02 (m/sec).

SEA SURFACE TEMPERATURE (DEG.C)

DEC 77




FIGURE 2.6 Monthly mean sea surface temperature is the mean of 12-hourly FNWC sea surface temperatures. Contour intervals are 1°C.

AIR TEMPERATURE (DEG. C) DEC 77




FIGURE 2.7 Monthly mean air temperature is the mean of 12-hourly FNWC air temperature. Contour intervals are 2°C.

700 MB HEIGHT (M) DEC 77




FIGURE 2.8 Monthly mean 700 mb height is the mean of 12-hourly FNWC 700 mb heights. Contour intervals are 25 meters.

VAPOR PRESSURE (MB)

DEC 77




FIGURE 2.9 Monthly mean surface vapor pressure is the mean of 12-hourly FNWC vapor pressure at 19.5 meters. Contour intervals are 1 mb.

SENSIBLE HEAT FLUX ( $10^{-4}$  CAL/CM<sup>2</sup> SEC)

DEC 77




FIGURE 2.10 Monthly mean sensible heat flux (ocean to atmosphere) is the mean of 12-hourly sensible heat flux calculated from FNWC air and sea temperature and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $1.0 \times 10^{-4}$  cal/cm<sup>2</sup> sec.




FIGURE 2. II Monthly mean latent heat flux (ocean to atmosphere) is the mean of 12-hourly latent heat flux calculated from FNWC sea temperature, vapor pressure and wind using a bulk formula. Iso-line of zero heat flux are plotted heavily, and contour intervals are  $0.5 \times 10^{-3}$  cal/cm $^2$  sec.




FIGURE 2.12 Monthly temperature anomalies ( $^{\circ}\text{C}$ ) contoured at fixed depths. Negative anomaly areas are hatched and isolines of zero anomaly are drawn heavily. The crosses mark the positions of the TRANSPAC XBT drops. Increments are in  $0.1^{\circ}\text{C}$  (prepared by N. White, SIO).






FIGURE 2.13 Monthly buoy tracks. Tracks are drawn from starting point at beginning of month to terminal point at end of month. Ratio of latitude to longitude is the same as in other figures.

WIND SPEED (M/SEC)

JAN 78




FIGURE 3.1 Absolute value of monthly mean vector wind velocities at 19.5 meters. Contour intervals are 1 m/sec.

WIND DIRECTION

JAN

78




FIGURE 3.2 Direction arrows representing directions of monthly mean wind vectors at 19.5 meters. Length of arrow shaft indicates wind speed in m/sec. (See scale above figure).

WIND STRESS (DYNES/CM\*\*2)

JAN 78




FIGURE 3.3 Monthly mean wind stress is the mean of 6-hourly wind stress at 10 meters calculated from FNWC wind data. Contour intervals are 0.2 dynes/cm².

CURL OF WIND STRESS (10\*\*-9 DYNES/CM\*\*3)

JAN 78




FIGURE 3.4 The vertical component of monthly mean wind stress curl is the mean of 6-hourly wind stress curl approximated by finite-differences from 6-hourly wind stresses at 10 meters. Isolines of zero curl are plotted heavily, and contour intervals are  $4.0 \times 10^{-9}$  dynes/cm³.

U STAR CUBED ((M/SEC)\*\*3)

JAN 78




FIGURE 3.5 Monthly mean wind shear stress velocity cubed,  $U_*^3$  is the mean of 6-hourly wind shear stress velocity cubed calculated from wind speed at 10 meters. Contour intervals are 0.02 (m/sec).

SEA SURFACE TEMPERATURE (DEG.C)

JAN 78




FIGURE 3.6 Monthly mean sea surface temperature is the mean of 12-hourly FNWC sea surface temperatures. Contour intervals are 1°C.

AIR TEMPERATURE (DEG. C) JAN 78




FIGURE 3.7 Monthly mean air temperature is the mean of 12-hourly FNWC air temperature. Contour intervals are 2°C.

700 MB HEIGHT (M) JAN 78




FIGURE 3.8 Monthly mean 700 mb height is the mean of 12-hourly FNWC 700 mb heights. Contour intervals are 25 meters.

VAPOR PRESSURE (MB) JAN 78




FIGURE 3.9 Monthly mean surface vapor pressure is the mean of 12-hourly FNCW vapor pressure at 19.5 meters. Contour intervals are 1 mb.

SENSIBLE HEAT FLUX ( $10^{-4}$  CAL/CM<sup>2</sup> SEC) JAN 78




FIGURE 3.10 Monthly mean sensible heat flux (ocean to atmosphere) is the mean of 12-hourly sensible heat flux calculated from FNCW air and sea temperature and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $1.0 \times 10^{-4}$  cal/cm<sup>2</sup> sec.




FIGURE 3.11 Monthly mean latent heat flux (ocean to atmosphere) is the mean of 12-hourly latent heat flux calculated from FNWC sea temperature, vapor pressure and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $0.5 \times 10^{-3}$  cal/cm<sup>2</sup> sec.

# OPTIMALLY INTERPOLATED TRANSPAC SST OBS. MAR. 1972




FIGURE 3.12 Monthly temperature anomalies (°C) contoured at fixed depths. Negative anomaly areas are hatched and isolines of zero anomaly are drawn heavily. The crosses mark the positions of the TRANSPAC SST drops. Interpolated are in 1°C (prepared by M. White, 1972).



# BUOY PLOT FOR JAN78




FIGURE 3.13 Monthly buoy tracks. Tracks are drawn from starting point at beginning of month to terminal point at end of month. Ratio of latitude to longitude is the same as in other figures.

WIND SPEED (M/SEC)

FEB 78




FIGURE 4.1 Absolute value of monthly mean vector wind velocities at 19.5 meters. Contour intervals are 1 m/sec.

WIND DIRECTION

FEB

78




FIGURE 4.2 Direction arrows representing directions of monthly mean wind vectors at 19.5 meters. Length of arrow shaft indicates wind speed in m/sec. (See scale above figure).




FIGURE 4.3 Monthly mean wind stress is the mean of 6-hourly wind stress at 10 meters calculated from FNWC wind data. Contour intervals are  $0.2 \text{ dynes/cm}^2$ .





FIGURE 4.4 The vertical component of monthly mean wind stress curl is the mean of 6-hourly wind stress curl approximated by finite-differences from 6-hourly wind stresses at 10 meters. Isolines of zero curl are plotted heavily, and contour intervals are  $4.0 \times 10^{10} \text{ dynes/cm}^3$ .

U STAR CUBED ((M/SEC)\*\*3)

FEB 78



AIR TEMPERATURE (DEG. C) FEB 78




FIGURE 4.7 Monthly mean air temperature is the mean of 12-hourly FNWC air temperature. Contour intervals are 2°C.

700 MB HEIGHT (M) FEB 78




FIGURE 4.8 Monthly mean 700 mb height is the mean of 12-hourly FNWC 700 mb heights. Contour intervals are 25 meters.




FIGURE 4.9 Monthly mean surface vapor pressure is the mean of 12-hourly FNWC vapor pressure at 19.5 meters. Contour intervals are 1 mb.




FIGURE 4.10 Monthly mean sensible heat flux (ocean to atmosphere) is the mean of 12-hourly sensible heat flux calculated from FNWC air and sea temperature and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $1.0 \times 10^{-4}$  cal/cm $\cdot$ 2 sec.




FIGURE #11 Monthly mean latent heat flux (ocean to atmosphere) is the mean of 12-hourly latent heat flux calculated from FNWC sea temperature, vapor pressure and wind using a bulk formula. Isofine of zero heat flux are plotted heavily, and contour intervals are  $0.5 \times 10^{-3}$  cal/cm $\cdot$ 2 sec.

# OPTIMALLY INTERPOLATED TRANSPAC XBT OBS-FEB 1978




FIGURE 4.12 Monthly temperature anomalies (°C) contoured at fixed depths. Negative anomaly areas are hatched and isoline of zero anomaly are drawn heavily. The crosses mark the positions of the TRANSPAC XBT drops. Increments are in .1°C (prepared by W. White, SIO).



# BUOY PLOT FOR FEB78




FIGURE 4.13 Monthly buoy tracks. Tracks are drawn from starting point at beginning of month to terminal point at end of month. Ratio of latitude to longitude is the same as in other figures.

WIND SPEED (M/SEC)

MAR 78




FIGURE 5.1 Absolute value of monthly mean vector wind velocities at 19.5 meters. Contour intervals are 1 m/sec.

WIND DIRECTION

MAR

78




FIGURE 5.2 Direction arrows representing directions of monthly mean wind vectors at 19.5 meters. Length of arrow shaft indicates wind speed in m/sec. (See scale above figure).

WIND STRESS (DYNES/CM\*\*2)

MAR 78




FIGURE 5.3 Monthly mean wind stress is the mean of 6-hourly wind stress at 10 meters calculated from FNWC wind data. Contour intervals are 0.2 dynes/cm².

CURL OF WIND STRESS (10\*\*-9 DYNES/CM\*\*3)

MAR 78




FIGURE 5.4 The vertical component of monthly mean wind stress curl is the mean of 6-hourly wind stress curl approximated by finite-differences from 6-hourly wind stresses at 10 meters. Isolines of zero curl are plotted heavily, and contour intervals are  $4.0 \times 10^{-9}$  dynes/cm³.

U STAR CUBED ((M/SEC)\*\*3)

MAR 78




FIGURE 5.5 Monthly mean wind shear stress velocity cubed,  $U_*^3$  is the mean of 6-hourly wind shear stress velocity cubed calculated from wind speed at 10 meters. Contour intervals are 0.02 (m/sec)<sup>3</sup>.

SEA SURFACE TEMPERATURE (DEG.C)

MAR 78




FIGURE 5.6 Monthly mean sea surface temperature is the mean of 12-hourly FNWC sea surface temperatures. Contour intervals are 1°C.

AIR TEMPERATURE (DEG. C) MAR 78




FIGURE 5.7 Monthly mean air temperature is the mean of 12-hourly FNWC air temperature. Contour intervals are 2°C.

700 MB HEIGHT (M) MAR 78




FIGURE 5.8 Monthly mean 700 mb height is the mean of 12-hourly FNWC 700 mb heights. Contour intervals are 25 meters.

VAPOR PRESSURE (MB) MAR 78




FIGURE 5.9 Monthly mean surface vapor pressure is the mean of 12-hourly FNWC vapor pressure at 19.5 meters. Contour intervals are 1 mb.

SENSIBLE HEAT FLUX ( $10^{-4}$  CAL/CM<sup>2</sup> SEC) MAR 78




FIGURE 5.10 Monthly mean sensible heat flux (ocean to atmosphere) is the mean of 12-hourly sensible heat flux calculated from FNWC air and sea temperature and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $1.0 \times 10^{-4}$  cal/cm<sup>2</sup> sec.




FIGURE 5.11 Monthly mean latent heat flux (ocean to atmosphere) is the mean of 12-hourly latent heat flux calculated from FNWC sea temperature, vapor pressure and wind using a bulk formula; isoline of zero heat flux are plotted heavily, and contour intervals are  $0.5 \times 10^{-3}$  cal/cm $^2$  sec.

# DIFFUSIVELY INTERPOLATED TRANSPAC XBT 055 MAR 1978




FIGURE 5.4 Monthly temperature anomalies (°C) contoured at fixed depths. Negative anomaly areas are hatched and isolines of zero anomaly are drawn heavily. The crosses mark the positions of the TRANSPAC XBT drops. Increments are in 1.0°C (prepared by M. Martin, CIO).






FIGURE 5.13 Monthly buoy tracks. Tracks are drawn from starting point at beginning of month to terminal point at end of month. Ratio of latitude to longitude is the same as in other figures.

WIND SPEED (M/SEC)

APR /8




FIGURE 6.1 Absolute value of monthly mean vector wind velocities at 19.5 meters. Contour intervals are 1 m/sec.

WIND DIRECTION

APR

78




FIGURE 6.2 Direction arrows representing directions of monthly mean wind vectors at 19.5 meters. Length of arrow shaft indicates wind speed in m/sec. (See scale above figure).




FIGURE 6.3 Monthly mean wind stress is the mean of 6-hourly wind stress at 10 meters calculated from FNWC wind data. Contour intervals are 0.2 dynes/cm<sup>2</sup>.




FIGURE 6.4 The vertical component of monthly mean wind stress curl is the mean of 6-hourly wind stress curl approximated by finite-differences from 6-hourly wind stresses at 10 meters. Isolines of zero curl are plotted heavily, and contour intervals are  $4.0 \times 10^{-9}$  dynes/cm<sup>3</sup>.

U STAR CUBED ((M/SEC)\*\*3)

APR 78




FIGURE 6.5 Monthly mean wind shear stress velocity cubed,  $U_*^3$  is the mean of 6-hourly wind shear stress velocity cubed calculated from wind speed at 10 meters. Contour intervals are 0.02 (m/sec).

SEA SURFACE TEMPERATURE (DEG.C)

APR 78




FIGURE 6.6 Monthly mean sea surface temperature is the mean of 12-hourly FNWC sea surface temperatures. Contour intervals are 1°C.

AIR TEMPERATURE (DEG. C)      APR    78




FIGURE 6.7 Monthly mean air temperature is the mean of 12-hourly FNWC air temperature. Contour intervals are 2°C.

700 MB HEIGHT (M)      APR    78




FIGURE 6.8 Monthly mean 700 mb height is the mean of 12-hourly FNWC 700 mb heights. Contour intervals are 25 meters.

VAPOR PRESSURE (MB)      APR    78




FIGURE 6.9 Monthly mean surface vapor pressure is the mean of 12-hourly FNWC vapor pressure at 19.5 meters. Contour intervals are 1 mb.

SENSIBLE HEAT FLUX ( $10^{-4}$  CAL/CM<sup>2</sup> SEC)      APR    78




FIGURE 6.10 Monthly mean sensible heat flux (ocean to atmosphere) is the mean of 12-hourly sensible heat flux calculated from FNWC air and sea temperature and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $1.0 \times 10^{-4}$  cal/cm<sup>2</sup> sec.




FIGURE 6.11 Monthly mean latent heat flux (ocean to atmosphere) is the mean of 12-hourly latent heat flux calculated from FNWC sea temperature, vapor pressure and wind using a bulk formula. Iso-line of zero heat flux are plotted heavily, and contour intervals are  $0.5 \times 10^{-3}$  cal/cm<sup>2</sup> sec.

# OPTIMALLY INTERPOLATED TRANSPAC XBT OBS. AREA 1978




FIGURE 6.12 Monthly temperature anomalies (°C) contoured at fixed depths. Negative anomaly areas are hatched and isolines of zero anomaly are drawn heavily. The crosses mark the positions of the TRANSPAC XBT drops. Increments are in 1°C (prepared by W. White, SIO).



# BUOY PLOT FOR APR 78




FIGURE 6.13 Monthly buoy tracks. Tracks are drawn from starting point at beginning of month to terminal point at end of month. Ratio of latitude to longitude is the same as in other figures.

WIND SPEED (M/SEC)

MAY 78




FIGURE 7.1 Absolute value of monthly mean vector wind velocities at 19.5 meters. Contour intervals are 1 m/sec.

WIND DIRECTION

MAY

78




FIGURE 7.2 Direction arrows representing directions of monthly mean wind vectors at 19.5 meters. Length of arrow shaft indicates wind speed in m/sec. (See scale above figure).

WIND STRESS (DYNES/CM\*\*2)

MAY 78




FIGURE 7.3 Monthly mean wind stress is the mean of 6-hourly wind stress at 10 meters calculated from FNCW wind data. Contour intervals are 0.2 dynes/cm².

CURL OF WIND STRESS ( $10^{-9}$  DYNES/CM\*\*3)

MAY 78




FIGURE 7.4 The vertical component of monthly mean wind stress curl is the mean of 6-hourly wind stress curl approximated by finite-differences from 6-hourly wind stresses at 10 meters. Isolines of zero curl are plotted heavily, and contour intervals are  $4.0 \times 10^{-9}$  dynes/cm³.

U STAR CUBED ((M/SEC)\*\*3)

MAY 78




FIGURE 7.5 Monthly mean wind shear stress velocity cubed,  $U_*^3$  is the mean of 6-hourly wind shear stress velocity cubed calculated from wind speed at 10 meters. Contour intervals are 0.02  $(m/sec)^3$ .

SEA SURFACE TEMPERATURE (DEG.C)

MAY 78




FIGURE 7.6 Monthly mean sea surface temperature is the mean of 12-hourly FNWC sea surface temperatures. Contour intervals are 1°C.

AIR TEMPERATURE (DEG. C) MAY 78




FIGURE 7.7 Monthly mean air temperature is the mean of 12-hourly FNWC air temperature. Contour intervals are 2°C.

700 MB HEIGHT (M) MAY 78




FIGURE 7.8 Monthly mean 700 mb height is the mean of 12-hourly FNWC 700 mb heights. Contour intervals are 25 meters.

VAPOR PRESSURE (MB) MAY 78




FIGURE 7.9 Monthly mean surface vapor pressure is the mean of 12-hourly FNWC vapor pressure at 19.5 meters. Contour intervals are 1 mb.

SENSIBLE HEAT FLUX ( $10^{-4}$  CAL/CM<sup>2</sup> SEC) MAY 78




FIGURE 7.10 Monthly mean sensible heat flux (ocean to atmosphere) is the mean of 12-hourly sensible heat flux calculated from FNWC air and sea temperature and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $1.0 \times 10^{-4}$  cal/cm<sup>2</sup> sec.

LATENT HEAT FLUX ( $10^{-3}$  CAL/CM<sup>2</sup> SEC)

MAY 78




FIGURE 7.11 Monthly mean latent heat flux (ocean to atmosphere) is the mean of 12-hourly latent heat flux calculated from FNWC sea temperature, vapor pressure and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $0.5 \times 10^{-3}$  cal/cm<sup>2</sup> sec.




FIGURE 7.12 Monthly temperature anomalies ( $^{\circ}\text{C}$ ) contoured at fixed depths. Negative anomaly areas are hatched and isoline of zero anomaly are drawn heavily. The crosses mark the positions of the TPANSPAC XBT drops. Increments are in  $0.1^{\circ}\text{C}$  (prepared by W. White, SIO).






FIGURE 7.13 Monthly buoy tracks. Tracks are drawn from starting point at beginning of month to terminal point at end of month. Ratio of latitude to longitude is the same as in other figures.

## APPENDIX. Analysis Procedure

Since some revisions have been made in the calculation of curl of wind stress, and since it has been some time since the publication of ADS Report Number 1, which contained a complete outline of the analysis procedure, this appendix is included in this report.

FNOC data were taken from 63 x 63 polar gridded fields of 6-hourly wind speed and direction, and 12-hourly 700 mb height, sea surface temperature, air temperature, and vapor pressure. The wind at a ten meter height above sea level was estimated from FNOC 19.5 meter winds by an iterative scheme using the neutral flux-profile relationships obtained by Businger et al. (1971). Wind stress was calculated from FNOC wind speed and direction using a bulk aerodynamic equation:

$$\tau = -\rho \bar{u}\bar{w} = \rho C_d U_{10}^2 \quad (\text{A.1})$$

where

$\tau$  = surface stress (dynes/cm<sup>2</sup>)

$\bar{u}\bar{w}$  = Reynolds stress, (m<sup>2</sup>sec<sup>-2</sup>)

$\rho$  = air density = 1.2923 x 10<sup>-3</sup> g/cm<sup>3</sup>

$U_{10}$  = wind speed at 10 m (cm/sec)

$C_d = 1.3 \times 10^{-3}$

The drag coefficient  $C_d$  was found by Smith and Banke (1975) for  $U_{10} = 10$  m/sec. Vertical component of wind stress curl was calculated on the FNOC northern hemispheric grid using polar projection mapping and finite difference approximations.

The vertical component of wind stress curl is,

$$\hat{r} \cdot \text{curl} \vec{\tau} = \frac{1}{R \sin \theta} \left[ \frac{\partial(\tau_\phi \sin \theta)}{\partial \theta} - \frac{\partial \tau_\theta}{\partial \phi} \right] \quad (\text{A.2})$$

where  $\theta$  and  $\phi$  are co-latitude and east longitude in polar coordinates, respectively, and  $\tau$  is wind stress,  $R$  is the Earth's radius, and  $\hat{r}$  is a unit radius vector.

A finite differencing method will be used to estimate this equation. However, since the data resides on a polar stereographic grid, a finite differencing scheme must be expressed in terms of polar stereographic coordinates, and then related to co-latitude and longitude using

the chain rule. The finite differences are:

$$\begin{aligned}\frac{\partial \tau_\theta}{\partial \zeta} &= \tau_\theta \left| \frac{\zeta_{i+1}}{\zeta_{i-1}} / (\zeta_{i+1} - \zeta_{i-1}) \right| \\ &= \tau_\theta \left| \frac{\zeta_{i+1}}{\zeta_{i-1}} / 2; i=1, 63 \right|\end{aligned}\quad (\text{A.3})$$

$$\begin{aligned}\frac{\partial \tau_\theta}{\partial \eta} &= \tau_\theta \left| \frac{\eta_{j+1}}{\eta_{j-1}} / (\eta_{j+1} - \eta_{j-1}) \right| \\ &= \tau_\theta \left| \frac{\eta_{j+1}}{\eta_{j-1}} / 2; j=1, 63 \right|\end{aligned}\quad (\text{A.4})$$

$$\begin{aligned}\frac{\partial (\tau_\phi \sin \zeta)}{\partial \zeta} &= \tau_\phi \sin \theta \left| \frac{\zeta_{i+1}}{\zeta_{i-1}} / (\zeta_{i+1} - \zeta_{i-1}) \right| \\ &= \tau_\phi \sin \theta \left| \frac{\zeta_{i+1}}{\zeta_{i-1}} / 2; i=1, 63 \right|\end{aligned}\quad (\text{A.5})$$

$$\begin{aligned}\frac{\partial (\tau_\phi \sin \theta)}{\partial \eta} &= \tau_\phi \sin \theta \left| \frac{\eta_{j+1}}{\eta_{j-1}} / (\eta_{j+1} - \eta_{j-1}) \right| \\ &= \tau_\phi \sin \theta \left| \frac{\eta_{j+1}}{\eta_{j-1}} / 2; j=1, 63 \right|\end{aligned}\quad (\text{A.6})$$

where  $\zeta(\theta, \phi)$  and  $\eta(\theta, \phi)$  are the abscissa and ordinate grid axes of the  $63 \times 63$  polar stereographic grid.

The partial derivatives may be calculated for the polar stereographic mapping equations:

$$\zeta = 32 + 31.205 \left( \frac{1 - \cos \theta}{1 + \cos \theta} \right)^{1/2} \cdot \sin(\phi + 80); i=1, 63, \quad (\text{A.7})$$

$$\eta = 32 - 31.205 \left( \frac{1 - \cos \theta}{1 + \cos \theta} \right)^{1/2} \cdot \cos(\phi + 80); j=1, 63, \quad (\text{A.8})$$

and the partial derivatives of  $\zeta$  and  $\eta$  with respect to  $\theta$  and  $\phi$  are:

$$\frac{\partial \zeta}{\partial \phi} = 31.205 \left( \frac{1 - \cos \theta}{1 + \cos \theta} \right)^{1/2} \cdot \cos(\phi + 80) \quad (\text{A.9})$$

$$\frac{\partial \eta}{\partial \phi} = 31.205 \left( \frac{1 - \cos \theta}{1 + \cos \theta} \right)^{1/2} \cdot \sin(\phi + 80) \quad (\text{A.10})$$

$$\frac{\partial \zeta}{\partial \theta} = 31.205 \left( \frac{1 - \cos \theta}{1 + \cos \theta} \right)^{-1/2} \cdot \frac{\sin \theta}{(1 + \cos \theta)^2} \sin(\phi + 80) \quad (\text{A.11})$$

$$\frac{\partial \eta}{\partial \theta} = -31.205 \left( \frac{1 - \cos \theta}{1 + \cos \theta} \right)^{-1/2} \cdot \frac{\sin \theta}{(1 + \cos \theta)^2} \cos(\phi + 80) \quad (\text{A.12})$$

The vertical component of the wind stress curl can then be estimated, using equations A.2 through A.12, as:

$$\begin{aligned}\hat{r} \cdot \text{curl} \vec{\tau} &= \frac{15.6025}{R \sin \theta_{ij}} \left( \frac{1 - \cos \theta_{ij}}{1 + \cos \theta_{ij}} \right)^{1/2} \cdot \left[ \tau_\phi \sin \theta \left| \frac{\zeta_{i+1}}{\zeta_{i-1}} \frac{\sin(\phi + 80)}{\sin \theta} \right|_{i,j} \right. \\ &\quad \left. - \tau_\phi \sin \theta \left| \frac{\eta_{j+1}}{\eta_{j-1}} \frac{\cos(\phi + 80)}{\sin \theta} \right|_{i,j} \right. \\ &\quad \left. - \tau_\theta \left| \frac{\zeta_{i+1}}{\zeta_{i-1}} \cos(\phi + 80) \right|_{i,j} - \tau_\theta \left| \frac{\eta_{j+1}}{\eta_{j-1}} \sin(\phi + 80) \right|_{i,j} \right]\end{aligned}\quad (\text{A.13})$$

Sensible heat flux ( $\text{cal}/\text{cm}^2\text{sec}$ ) was obtained using an empirical bulk formula (Friehe and Schmitt, 1976):

$$S.H.F. = \rho C_p [0.2 + C_h \cdot U_{10} \cdot (T_s - T_a)] \quad (\text{A.14})$$

where

$$C_p = 0.24 \text{ cal}/\text{gm}^\circ\text{C}$$

$$C_h = 0.91 \times 10^{-3}$$

$$T_s = \text{sea surface temperature}$$

$$T_a = (10 \text{ meter}) \text{ air temperature}$$

The latent heat flux ( $\text{cal}/\text{cm}^2\text{sec}$ ) was also calculated from a bulk formula (Friehe and Schmitt, 1976):

$$L.H.F. = L \cdot C_e \cdot U_{10} [Q_{\text{sea}} - Q] \quad (\text{A.15})$$

where

$$L = 595 \text{ cal}/\text{gm} \text{ (heat of evaporation)}$$

$$C_e = 1.32 \times 10^{-3}$$

$$Q = 0.75 \cdot [EAIR (\text{gm} \cdot \text{m}^{-3}) \equiv \text{FNOG vapor pressure}]$$

$$Q_{\text{sea}} = 1.667 \times 10^{-7} \cdot [e^{17.19 \cdot [(T_s + 273) / 273]}]$$

Stress, stress-curl and  $U^3$  fields were calculated at 00z, 06z, 12z ; heat fluxes were calculated for 00z and 12z every day. These calculated fields were then averaged to obtain daily, 5-daily, and monthly vector means of wind speed and direction. Monthly means were contoured over the region of the North Pacific from 120E to 230E and 20N to 60N.

MANADATORY DISTRIBUTION LIST  
FOR UNCLASSIFIED TECHNICAL REPORTS, REPRINTS & FINAL REPORTS  
PUBLISHED BY OCEANOGRAPHIC CONTRACTORS  
OF THE OCEAN SCIENCE AND TECHNOLOGY DIVISION  
OF THE OFFICE OF NAVAL RESEARCH  
(Revised July 1978)

Department of Defense

Office of the Secretary of Defense (3)  
Assistant Director of Defense Research  
& Engineering  
Washington, D.C. 20301

Naval Research Laboratory (6)  
Library, Code 2620  
Washington, D.C. 20375

U.S. Naval Oceanographic Office  
Library, Code 8170  
NSTL Station  
Bay St. Louis, MS 39529

Navy

Office of Naval Research (3)  
Code 460  
Arlington, VA 22217

Office of Naval Research  
Code 480  
Arlington, VA 22217

Office of Naval Research  
Code 102 B  
Arlington, VA 22217

Office of Naval Research (6)  
Code 102 DI  
Arlington, VA 22217

Office of Naval Research  
Commanding Officer  
1030 East Green Street  
Pasadena, CA 91101

Naval Ocean Research & Development  
Activity  
NORDA, Code 300  
NSTL Station  
Bay St. Louis, MS 39529

Other Government Agencies

Defense Documentation Center (12)  
Cameron Station  
Alexandria, VA 22314

National Oceanic & Atmospheric  
Administration  
National Oceanographic Data Center  
Washington Navy Yard  
Rockville, MD 20852